Computer Graphics (Fall 2005)

COMS 4160, Lecture 5: Viewing http://www.cs.columbia.edu/~cs4160

To Do


- Questions/concerns about assignment 1?
- Remember it is due Thu. Ask me or TA if any problems.

Motivation

- We have seen transforms (between coord systems)
- But all that is in 3D
- We still need to make a 2D picture
- Project 3D to 2D. How do we do this?
- This lecture is about viewing transformations

Demo (Projection Tutorial)

- Nate Robbins OpenGL tutors
- Projection exe
- Download others



What we've seen so far

- Transforms (translation, rotation, scale) as 4x4 homogeneous matrices
- Last row always 0 0 0 1. Last w component always 1
- For viewing (perspective), we will use that last row and w component no longer 1 (must divide by it)

Outline

- Orthographic projection (simpler)
- Perspective projection, basic idea
- Derivation of gluPerspective (handout: glFrustum)
- Brief discussion of nonlinear mapping in z

Not well covered in textbook chapter 7. We follow section 3.5 of real-time rendering most closely. Handouts on this will be given out.

Projections

- To lower dimensional space (here 3D -> 2D)
- Preserve straight lines
- Trivial example: Drop one coordinate (Orthographic)


Orthographic Projection Characteristic: Parallel lines remain parallel Useful for technical drawings etc.

Fig 7.1 in text

Perspective

Example


Simply project onto xy plane, drop z coordinate



In general


- We have a cuboid that we want to map to the normalized or square cube from [-1, +1] in all axes
- We have parameters of cuboid (l,r; t,b; n,f)

Orthographic




Orthographic Matrix

- First center cuboid by translating
- Then scale into unit cube



Caveats

- Looking down -z, f and n are negative (n > f)
- OpenGL convention: positive n, f, negate internally



Transformation Matrix

Scale

Translation (centering)

$$M = \begin{pmatrix} \frac{2}{r-l} & 0 & 0 & 0\\ 0 & \frac{2}{t-b} & 0 & 0\\ 0 & 0 & \frac{2}{f-n} & 0\\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & -\frac{l+r}{2}\\ 0 & 1 & 0 & -\frac{t+b}{2}\\ 0 & 0 & 1 & -\frac{f+n}{2}\\ 0 & 0 & 0 & 1 \end{pmatrix}$$

Final Result


$$M = \begin{pmatrix} \frac{2}{r-l} & 0 & 0 & -\frac{r+l}{r-l} \\ 0 & \frac{2}{t-b} & 0 & -\frac{t+b}{t-b} \\ 0 & 0 & \frac{2}{f-n} & -\frac{f+n}{f-n} \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad glOrtho = \begin{pmatrix} \frac{2}{r-l} & 0 & 0 & -\frac{r+l}{r-l} \\ 0 & \frac{2}{t-b} & 0 & -\frac{t+b}{t-b} \\ 0 & 0 & \frac{2}{f-n} & -\frac{f+n}{f-n} \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

Outline


- Orthographic projection (simpler)
- Perspective projection, basic idea
- Derivation of gluPerspective (handout: glFrustum)
- Brief discussion of nonlinear mapping in z

Perspective Projection

- Most common computer graphics, art, visual system
- Further objects are smaller (size, inverse distance)
- Parallel lines not parallel; converge to single point



Overhead View of Our Screen



Looks like we've got some nice similar triangles here?

$$\frac{x}{z} = \frac{x'}{d} \Rightarrow x' = \frac{d * x}{z}$$
 $\frac{y}{z} = \frac{y'}{d} \Rightarrow y' = \frac{d * y}{z}$

In Matrices

- Note negation of z coord (focal plane –d)
- Only) last row affected (no longer 0 0 0 1)
- w coord will no longer = 1. Must divide at end


$$P = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & -\frac{1}{d} & 0 \end{pmatrix}$$


Verify


$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & -\frac{1}{d} & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \\ 1 \end{pmatrix} = ? \qquad \begin{pmatrix} x \\ y \\ z \\ -\frac{1}{d} \end{pmatrix} = \begin{pmatrix} -\frac{d*x}{z} \\ -\frac{d*y}{z} \\ -d \\ 1 \end{pmatrix}$$

Outline

- Orthographic projection (simpler)
- Perspective projection, basic idea
- Derivation of gluPerspective (handout: glFrustum)
- Brief discussion of nonlinear mapping in z








gluPerspective

- gluPerspective(fovy, aspect, zNear > 0, zFar > 0)
- Fovy, aspect control fov in x, y directions
- zNear, zFar control viewing frustum

Overhead View of Our Screen



$$\theta = \frac{fovy}{2} \qquad d = \cot \theta$$

In Matrices

Simplest form:
$$P = \begin{pmatrix} \frac{1}{aspect} & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & -\frac{1}{d} & 0 \end{pmatrix}$$

- Aspect ratio taken into account
- Homogeneous, simpler to multiply through by d
- Must map z values based on near, far planes (not yet)

In Matrices

$$P = \begin{pmatrix} \frac{1}{aspect} & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & -\frac{1}{d} & 0 \end{pmatrix} \rightarrow \begin{pmatrix} \frac{d}{aspect} & 0 & 0 & 0 \\ 0 & d & 0 & 0 \\ 0 & 0 & A & B \\ 0 & 0 & -1 & 0 \end{pmatrix}$$

A and B selected to map n and f to -1, +1 respectively

Z mapping derivation

$$\begin{pmatrix} A & B \\ -1 & 0 \end{pmatrix} \begin{pmatrix} z \\ 1 \end{pmatrix} = ? \qquad \begin{pmatrix} Az + B \\ -z \end{pmatrix} = -A - \frac{B}{z}$$

Simultaneous equations?

$$-A + \frac{B}{n} = -1$$

$$-A + \frac{B}{f} = +1$$

$$A = -\frac{f+n}{f-n}$$

$$B = -\frac{2fn}{f-n}$$

Outline

- Orthographic projection (simpler)
- Perspective projection, basic idea
- Derivation of gluPerspective (handout: glFrustum)
- Brief discussion of nonlinear mapping in z

Mapping of Z is nonlinear

$$\begin{pmatrix} Az + B \\ -z \end{pmatrix} = -A - \frac{B}{z}$$

- Many mappings proposed: all have nonlinearities
- Advantage: handles range of depths (10cm 100m)
- Disadvantage: depth resolution not uniform
- More close to near plane, less further away
- Common mistake: set near = 0, far = infty. Don't do this. Can't set near = 0; lose depth resolution.
- We discuss this more in review session

