

Computer Graphics (Fall 2008)

COMS 4160, Lecture 3: Transformations 1



<http://www.cs.columbia.edu/~cs4160>

To Do

- Start (thinking about) assignment 1
 - Much of information you need is in this lecture (slides)
 - Ask TA NOW if compilation problems, visual C++ etc.
 - Not that much coding [solution is approx. 20 lines, but you may need more to implement basic matrix/vector math], but some thinking and debugging likely involved
- Specifics of HW 1
 - Axis-angle rotation and gluLookAt most useful (essential?). These are not covered in text (look at slides).
 - You probably only need final results, but try understanding derivations.
- Problems in text help understanding material. Usually, we have review sessions per unit, but this one before midterm

Course Outline

- 3D Graphics Pipeline



Course Outline

- 3D Graphics Pipeline



Unit 1: Transformations
Resizing and placing objects in the world. Creating perspective images.
Weeks 1 and 2
Ass 1 due Sep 25 ([Demo](#))

Motivation

- Many different coordinate systems in graphics
 - World, model, body, arms, ...
- To relate them, we must transform between them
- Also, for modeling objects. I have a teapot, but
 - Want to place it at correct location in the world
 - Want to view it from different angles (HW 1)
 - Want to scale it to make it bigger or smaller

Motivation

- Many different coordinate systems in graphics
 - World, model, body, arms, ...
- To relate them, we must transform between them
- Also, for modeling objects. I have a teapot, but
 - Want to place it at correct location in the world
 - Want to view it from different angles (HW 1)
 - Want to scale it to make it bigger or smaller
- This unit is about the math for doing all these things
 - Represent transformations using matrices and matrix-vector multiplications.
- Demo:** HW 1, applet [transformation_game.jar](#)

General Idea

- Object in model coordinates
 - Transform into world coordinates
 - Represent points on object as vectors
 - Multiply by matrices
 - Demos with applet
- Chapter 6 in text. We cover most of it essentially as in the book. Worthwhile (but not essential) to read whole chapter

Outline

- 2D transformations: rotation, scale, shear
- Composing transforms
- 3D rotations
- Translation: Homogeneous Coordinates (next time)
- Transforming Normals (next time)

(Nonuniform) Scale

$$Scale(s_x, s_y) = \begin{pmatrix} s_x & 0 \\ 0 & s_y \end{pmatrix} \quad S^{-1} = \begin{pmatrix} s_x^{-1} & 0 \\ 0 & s_y^{-1} \end{pmatrix}$$

$$\begin{pmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & s_z \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} s_x x \\ s_y y \\ s_z z \end{pmatrix}$$

[transformation_game.jar](#)

Shear

$$Shear = \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix} \quad S^{-1} = \begin{pmatrix} 1 & -a \\ 0 & 1 \end{pmatrix}$$



Rotations

2D simple, 3D complicated. [Derivation? Examples?]

$$2D? \quad \begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

- Linear $R(X+Y)=R(X)+R(Y)$
- Commutative

[transformation_game.jar](#)

Outline

- 2D transformations: rotation, scale, shear
- Composing transforms
- 3D rotations
- Translation: Homogeneous Coordinates
- Transforming Normals

Composing Transforms

- Often want to combine transforms
- E.g. first scale by 2, then rotate by 45 degrees
- Advantage of matrix formulation: All still a matrix
- Not commutative!! Order matters

E.g. Composing rotations, scales

$$x_3 = Rx_2 \quad x_2 = Sx_1$$

$$x_3 = R(Sx_1) = (RS)x_1$$

$$x_3 \neq SRx_1$$

[transformation_game.jar](#)

Inverting Composite Transforms

- Say I want to invert a combination of 3 transforms
- Option 1: Find composite matrix, invert
- Option 2: Invert each transform *and swap order*
- Obvious from properties of matrices

$$M = M_1 M_2 M_3$$

$$M^{-1} = M_3^{-1} M_2^{-1} M_1^{-1}$$

$$M^{-1}M = M_3^{-1}(M_2^{-1}(M_1^{-1}M_1)M_2)M_3$$

[transformation_game.jar](#)

Outline

- 2D transformations: rotation, scale, shear
- Composing transforms
- 3D rotations
- Translation: Homogeneous Coordinates
- Transforming Normals

Rotations

Review of 2D case

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

- Orthogonal?, $R^T R = I$

Rotations in 3D

- Rotations about coordinate axes simple

$$R_z = \begin{pmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad R_y = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \theta & -\sin \theta \\ 0 & \sin \theta & \cos \theta \end{pmatrix}$$

$$R_x = \begin{pmatrix} \cos \theta & 0 & \sin \theta \\ 0 & 1 & 0 \\ -\sin \theta & 0 & \cos \theta \end{pmatrix}$$

- Always linear, orthogonal $R^T R = I$
 - Rows/cols orthonormal $R(X+Y) = R(X) + R(Y)$

Axis-Angle: Putting it together

$$(b \setminus a)_{ROT} = (I_{3 \times 3} \cos \theta - aa^T \cos \theta)b + (A^* \sin \theta)b$$
$$(b \rightarrow a)_{ROT} = (aa^T)b$$

$$R(a, \theta) = I_{3 \times 3} \cos \theta + aa^T (1 - \cos \theta) + A^* \sin \theta$$

$$R(a, \theta) = \cos \theta \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + (1 - \cos \theta) \begin{pmatrix} x^2 & xy & xz \\ xy & y^2 & yz \\ xz & yz & z^2 \end{pmatrix} + \sin \theta \begin{pmatrix} 0 & -z & y \\ z & 0 & -x \\ -y & x & 0 \end{pmatrix}$$

(x y z) are cartesian components of a